Procedure Instruction Sheet for MIDWEST ENDOSCOPY CENTER, LLC www.sgihealth.com

DINESH JAIN, M.D. SCOTT BERGER, M.D. SUSHAMA GUNDLAPALLI, M.D. DARREN KASTIN, M.D. GONZALO PANDOLFI, M.D. SHIVANI KIRILUK. D.O. PRAVEEN METTU, M.D. ADITYA DHOLAKIA, D.O.

1243 Rickert Dr. Naperville, IL 60540

PEG 3350 PREPARATION

Telephone 630-527-6450 Fax 630-527-6456

SAME DAY BOWEL PREPARATION (FOR AFTERNOON PROCEDURES)

Your doctor has ordered a colonoscopy, which is an examination of the entire intestine (colon). This procedure takes about 20-30 minutes. A flexible tube (colonoscope) is passed into the rectum then gently advanced through the large intestine, examining the inner lining.

<u>10 days</u> before procedure	<u>5 days</u> before your procedure	The day before your Procedure. <u>STAY ON CLEAR</u> <u>LIQUID DIET</u> <u>THE ENTIRE DAY</u>	The morning of your procedure Start at: A.M. And finish within 4 hours.
Purchase PEG 3350 from your local pharmacy Avoid any foods with OLESTRA (found in low fat snack foods) Avoid popcorn, nuts or seeds.	Contact a relative or friend to arrange for transportation from the clinic after your procedure. BECAUSE YOU WILL BE SEDATED YOU MUST HAVE SOMEONE AVAILABLE TO TAKE YOU HOME YOU MAY NOT TAKE A TAXI HOME UNLESS YOU HAVE SOMEONE, OTHER THAN THE TAXI DRIVER, TO BE YOUR RESPONSIBLE ADULT.	See Clear liquid diet instructions.	Add drinking water to PEG 3350 to the fill line (4 liters) Mix thoroughly and chill. Patients may add Crystal Light® Lemonade mix to improve the taste. This is <u>optional</u> and <u>only the</u> <u>yellow lemonade</u> mix is permitted YOUR PREP MUST BE COMPLETED 4 HOURS PRIOR TO YOUR PROCEDUR DO NOT EAT ANYTHING! Remember to wear comfortable clothing Women please avoid high heels

Clear liquids mean anything that is clear in consistency: color is not as important as consistency. Drink juices such as apple, white grape or white cranberry. Avoid thick and pulpy juices. Bouillon or clear broth is fine. You can have gelatin without fruit. Avoid anything with red or orange artificial coloring. Soda, coffee and tea are fine. DO NOT drink milk or dairy products.

PLEASE SEE REVERSE SIDE **MIDWEST ENDOSCOPY CENTER, LLC**

www.sgihealth.com

If for any reason you are unable to complete your preparation of PEG 3350, please call our office and the answering service will page the doctor on call for you.

PATIENTS MAY HAVE CLEAR LIQUIDS UP TO 4 HOURS PRIOR TO PROCEDURE.

DAY OF EXAM

IF YOU ARE TAKING BLOOD THINNERS SUCH AS COUMADIN OR THE EQUIVALENT, OR DIABETIC MEDICATIONS, PLEASE INFORM THE DOCTOR, AS YOU WILL NEED SPECIAL **INSTRUCTIONS.**

- If you are taking medicine for high blood pressure, asthma, or heart disease you will want to take these as usual with a sip of water. All medications must be taken 2 hours prior to your procedure.
- All other medications must be reviewed with the doctor.
- Should you consume more than just a sip of water with your medications the day of your • procedure there will be a possibility of your procedure being cancelled.
- If polyps are removed from your colon (or if biopsies are taken) you may be told not to perform • strenuous physical activities for up to 2 weeks.
- If polyps are removed (or if biopsies are taken) you should not travel by airplane for 2 weeks. You ٠ should not travel overseas for 2 weeks.
- Please do not wear any facial make-up, moisturizers, body lotions or oils the day of your procedure.

Please call the office (630) 527-6450 if you have any questions regarding the instructions.

Please be aware not all preps have the same effect on everyone. Following the above instructions is not a guarantee that you will be properly prepared for your colonoscopy. There is a possibility that you will need to be rescheduled at a later date with additional instructions if your prep is inadequate.

Your procedure start time is not a guarantee.

Patient Name:DOB:/is scheduled for a Colonoscopy on:

Date: ______ Procedure Start Time: _____

PLEASE ARRIVE 30 MINUTES PRIOR TO YOUR START TIME. IF YOU ARRIVE EARLIER, YOU MAY BE WAITING LONGER.

Because you will be sedated, you will need a driver to bring you to and from your appointment. This driver MUST be with you at check in so we may verify you have a responsible adult bringing you home. We will be unable to go forward with your procedure if your driver is not with you at time of check in.

Report to: MIDWEST ENDOSCOPY CENTER, LLC 1243 Rickert Dr Naperville, IL 60540 630-527-6450 Special Instructions: